



(11)

EP 1 254 237 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention of the grant of the patent:
07.04.2010 Bulletin 2010/14

(51) Int Cl.:
C12N 15/53 (2006.01) **C12N 9/06 (2006.01)**
C12N 9/96 (2006.01) **A61K 47/48 (2006.01)**
A61K 38/44 (2006.01)

(21) Application number: **01923265.1**(86) International application number:
PCT/US2001/040069(22) Date of filing: **07.02.2001**(87) International publication number:
WO 2001/059078 (16.08.2001 Gazette 2001/33)

(54) AGGREGATE-FREE URATE OXIDASE FOR PREPARATION OF NON-IMMUNOGENIC POLYMER CONJUGATES

AGGREGAT-FREIE HARNSÄUREOXIDASE ZUR HERSTELLUNG VON NICHT-IMMUNOGENEN POLYMER-KONJUGATEN

URATE OXIDASE EXEMPTÉE D'AGRÉGATS DESTINÉE A LA PRÉPARATION DE POLYMÈRES CONJUGUÉS NON-IMMUNOGÈNES

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU MC NL PT SE TR

(30) Priority: **10.02.2000 US 501730**

(43) Date of publication of application:
06.11.2002 Bulletin 2002/45

(73) Proprietors:

- Mountain View Pharmaceuticals, Inc.
Menlo Park, CA 94025-1813 (US)
- Duke University
Durham, NC 27705 (US)

(72) Inventors:

- SHERMAN, Merry, R.
San Carlos, CA 94070 (US)
- SAIFER, Mark, G., P.
San Carlos, CA 94070 (US)
- WILLIAMS, L., David
Fremont, CA 94536 (US)
- HERSHFIELD, Michael, S.
Durham, Nort Carolina 27707 (US)
- KELLY, Susan, J.
Chapel Hill, North Carolina 27516 (US)

(74) Representative: **Evenson, Jane Harriet et al
Mathys & Squire LLP
120 Holborn
London EC1N 2SQ (GB)**

(56) References cited:
WO-A-00/07629

- MONTALBINI P ET AL: "URICASE FROM LEAVES: ITS PURIFICATION AND CHARACTERIZATION FROM THREE DIFFERENT HIGHER PLANTS" PLANTA, SPRINGER VERLAG, DE, vol. 202, no. 3, 8 January 1997 (1997-01-08), pages 277-283, XP000881523 ISSN: 0032-0935
- SUZUKI H ET AL: "SOYBEAN NODULE-SPECIFIC URICASE NODULIN 35 IS EXPRESSED AND ASSEMBLED INTO A FUNCTIONAL TETRAMERIC HOLOENZYME IN ESCHERICHIA-COLI" PLANT PHYSIOLOGY, AMERICAN SOCIETY OF PLANT PHYSIOLOGISTS, ROCKVILLE, MD, US, vol. 95, 1991, pages 384-389, XP002125213 ISSN: 0032-0889
- OSMAN A M ET AL: "Liver uricase in Camelus dromedarius: purification and properties" COMPARATIVE BIOCHEMISTRY AND PHYSIOLOGY. B. COMPARATIVE BIOCHEMISTRY, PERGAMON PRESS, LONDON, GB, vol. 94B, no. 3, 1989, pages 469-474, XP002125210 ISSN: 0305-0491

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

- YASUDA Y ET AL: "Biochemical and biopharmaceutical properties of macromolecular conjugates of uricase with dextran and polyethylene glycol" CHEMICAL AND PHARMACEUTICAL BULLETIN, PHARMACEUTICAL SOCIETY OF JAPAN. TOKYO, JP, vol. 38, no. 7, July 1990 (1990-07), pages 2053-2056, XP002125209 ISSN: 0009-2363
- DAVIS S ET AL: "HYPOURICAEMIC EFFECT OF POLYETHYLENEGLYCOL MODIFIED URATE OXIDASE" LANCET THE, LANCET LIMITED. LONDON, GB, 8 August 1981 (1981-08-08), pages 281-283, XP000577404 ISSN: 0140-6736
- SAVOCA K V ET AL: "Induction of tolerance in mice by uricase and monomethoxypolyethylene glycol-modified uricase" INTERNATIONAL ARCHIVES OF ALLERGY AND APPLIED IMMUNOLOGY, BASEL, CH, vol. 75, 1984, pages 58-67, XP002125212 ISSN: 0020-5915
- CHEN R H-L ET AL: "PROPERTIES OF TWO URATE OXIDASES MODIFIED BY THE COVALENT ATTACHMENT OF POLY(ETHYLENE GLYCOL)" BBA ENZYMOLOGY, ELSEVIER, AMSTERDAM, NL, vol. 660, no. 2, 13 August 1981 (1981-08-13), pages 293-298, XP000577602 ISSN: 0005-2744
- CHUA C C ET AL: "Use of polyethylene glycol-modified uricase (PEG - uricase) to treat hyperuricemia in a patient with non-Hodgkin lymphoma" ANNALS OF INTERNAL MEDICINE, NEW YORK, NY, US, 15 July 1988 (1988-07-15), pages 114-117, XP002125211 ISSN: 0003-4819
- BURNHAM N L: "POLYMERS FOR DELIVERING PEPTIDES AND PROTEINS" AMERICAN JOURNAL OF HOSPITAL PHARMACY, AMERICAN SOCIETY OF HOSPITAL PHARMACISTS, BETHESDA, MD, US, vol. 51, no. 2, 15 January 1994 (1994-01-15), pages 210-218, XP000783643 ISSN: 0002-9289
- TSUJI J.-I. ET AL: "Studies on antigenicity of the polyethylene glycol (PEG)-modified uricase" INTERNATIONAL JOURNAL OF IMMUNOPHARMACOLOGY, vol. 7, no. 5, 1 January 1985 (1985-01-01), pages 725-730, XP023812682 ELMSFORD, NY, US
- BRAUN A. ET AL: "Protein aggregates seem to play a key role among the parameters influencing the antigenicity of interferon alpha (IFN-alpha) in normal and transgenic mice" PHARMACEUTICAL RESEARCH, vol. 14, no. 10, 1 January 1997 (1997-01-01), pages 1472-1478, XP002492311 NEW YORK, NY, US

Remarks:

The file contains technical information submitted after the application was filed and not included in this specification

DescriptionStatement of Government Rights in the Invention

5 [0001] A portion of the research described in this application was made with support from the U.S.-Israel Binational Industrial Research and Development Foundation. Accordingly, the U.S. Government may have certain rights in the invention.

Background of the InventionField of the Invention

10 [0002] The present invention relates to purification and chemical modification of proteins to prolong their circulating lifetimes and reduce their immunogenicity. More specifically, the invention relates to the removal of aggregates larger than octamers from urate oxidases (uricases) prior to conjugation of poly(ethylene glycols) or poly(ethylene oxides). This substantially eliminates uricase immunogenicity without compromising its uricolytic activity.

Description of the Related Art

20 [0003] Statements contained in this background section do not constitute an admission of prior art, but instead reflect the inventors' own subjective comments on and interpretations of the state of the art at the time the invention was made. These interpretations may include personal, heretofore undisclosed, insights of the inventors, which insights were not themselves part of the prior art.

25 [0004] Urate oxidases (uricases; E.C. 1.7.3.3) are enzymes that catalyze the oxidation of uric acid to a more soluble product, allantoin, a purine metabolite that is more readily excreted. Humans do not produce enzymatically active uricase, as a result of several mutations in the gene for uricase acquired during the evolution of higher primates. Wu, X, et al., (1992) J Mol Evol 34:78-84. As a consequence, in susceptible individuals, excessive concentrations of uric acid in the blood (hyperuricemia) and in the urine (hyperuricosuria) can lead to painful arthritis (gout), disfiguring urate deposits (tophi) and renal failure. In some affected individuals, available drugs such as allopurinol (an inhibitor of uric acid synthesis) produce treatment-limiting adverse effects or do not relieve these conditions adequately. Hande, KR, et al., (1984) Am J Med 76:47-56; Fam, AG, (1990) Baillière's Clin Rheumatol 4:177-192. Injections of uricase can decrease hyperuricemia and hyperuricosuria, at least transiently. Since uricase is a foreign protein in humans, however, even the first injection of the unmodified protein from *Aspergillus flavus* has induced anaphylactic reactions in several percent of treated patients (Pui, C-H, et al., (1997) Leukemia 11:1813-1816), and immunologic responses limit its utility for chronic or intermittent treatment. Donadio, D, et al., (1981) Nouv Presse Méd 10:711-712; Leaustic, M, et al., (1983) Rev Rhum Mal Osteoartic 50:553-554.

30 [0005] U.S. Patent Application Serial No. 09/370,084 and published International Application No. PCT/US99/17514, disclose poly (ethylene glycol)-urate oxidase (PEG-uricase) that retains at least about 75% of the uricolytic activity of unconjugated uricase and has substantially reduced immunogenicity. In one such purified uricase, each subunit is covalently linked to an average of 2 to 10 sands of PEG, wherein each molecule of PEG may have a molecular weight between about 5 kDa and 100 kDa.

35 [0006] The aggregation of proteins is known to increase their immunogenicity. This understanding has contributed to the development of methods for intentionally aggregating proteins by treatments such as thermal denaturation and cross-linking by exposure to glutaraldehyde prior to use in the preparation of vaccines or for immunization of animals to produce antisera.

40 [0007] Unintentional aggregation of proteins has also been recognized as contributing to immunization or sensitization during clinical use of therapeutic proteins, e.g. for human gamma globulin (Henney et al. (1968) N. Engl. J. Med 278: 2244-2246) and for human growth hormone (Moore et al. (1980) J. Clin Endocrinol. Metab. 51:691-697). The contribution of aggregates to the immunogenicity of human interferon alpha has been demonstrated in BALB/c mice (Braun et al. (1997) Pharm. Res. 14:1472-1478) and an enzyme-linked immunosorbent assay (ELISA) has been developed for their measurement (Braun et al. (1997) Pharm. Res. 14:1394-1400).

45 [0008] In contrast to the known effects of aggregation on the immunogenicity of proteins, there are not reports of the effect of aggregation on the immunogenicity of proteins conjugated to poly(alkylene glycols) such as PEG. There is a need for poly(alkylene glycol)-uricase conjugates that substantially eliminate uricase immunogenicity without compromising its uricolytic activity. The present invention provides such compositions.

Summary of the Invention

[0009] Conjugation of proteins with poly(alkylene glycols), especially PEG, produces conjugates with reduced immunogenicity and increased persistence in the bloodstream. In attempting to produce substantially non-immunogenic conjugates of uricase that retain substantially all of the uricolytic activity of the unmodified uricase preparation, it was discovered that traces of large aggregates of uricase in the starting material were surprisingly effective at provoking both antibody formation and accelerated clearance from the circulation, both of which are deleterious, after repeated injections of PEG conjugates prepared from uricase containing such aggregates. Surprisingly, the present inventors found that the increased immunogenicity and accelerated clearance were *not* due to the presence of well-defined, moderate-sized aggregates of the uricase subunit that are larger than the native tetramer, e.g. aggregates containing eight subunits (octamers). The octameric form of uricase is present at sufficiently high concentrations in most preparations of uricase to be detectable by its absorbance of UV light, e.g. at 214 nm or 276 nm, or by its contribution to the refractive index or other measurements of protein concentration. Nevertheless, the octamers themselves were found to contribute minimally to the immunogenicity and accelerated clearance of PEG-uricase conjugates, in contrast with the much smaller quantities of the much larger aggregates that are undetectable by UV absorbance under the conditions tested but are readily detected by static (Raleigh) or dynamic light scattering. Therefore, the removal of such traces of very large aggregates prior to conjugation with PEG was found to decrease the immunogenicity and the accelerated clearance of the resultant PEG-uricase conjugates to a surprising extent.

[0010] One embodiment of the present invention is a uricase conjugate comprising a purified urate oxidase (uricase) conjugated to poly(ethylene glycol) or poly(ethylene oxide), wherein said uricase contains purified tetrameric and octameric forms of the uricase preparable by removing aggregates larger than octamers, wherein said uricase contains no more than about 2% of aggregates larger than octamers. Preferably, the uricase is mammalian uricase. More preferably, the uricase is porcine liver, bovine liver or ovine liver uricase. In one aspect of this preferred embodiment, the uricase is recombinant. In another aspect of this preferred embodiment, the uricase has substantially the sequence of porcine, bovine, ovine or baboon liver uricase. Advantageously, the uricase is chimeric. Preferably, the uricase is PKS uricase. In another aspect of this preferred embodiment, the uricase has substantially the sequence of baboon liver uricase in which tyrosine 97 has been replaced by histidine. Preferably, the uricase comprises an amino terminus and a carboxy terminus, and wherein the uricase is truncated at one terminus or both termini. Advantageously, the uricase is a fungal or microbial uricase. Preferably, the fungal or microbial uricase is isolated from *Aspergillus flavus*, *Arthrobacter globiformis*, *Bacillus* sp. or *Candida utilis*, or is a recombinant enzyme having substantially the sequence of one of said uricases. Alternatively, the uricase is an invertebrate uricase. Preferably, the invertebrate uricase is isolated from *Drosophila melanogaster* or *Drosophila pseudoobscura*, or is a recombinant enzyme having substantially the sequence of one of said uricases. In another aspect of this preferred embodiment, the uricase is a plant uricase. Preferably, the plant uricase is isolated from root nodules of *Glycine max* or is a recombinant enzyme having substantially the sequence of the uricase.

[0011] In one aspect of this preferred embodiment, the uricase described above is conjugated to poly(ethylene glycol) or poly(ethylene oxide), under conditions such that the uricase in the conjugate is substantially free of aggregates larger than octamers. Preferably, the uricase is conjugated to poly(ethylene glycol) or poly(ethylene oxide) via a urethane (carbamate), secondary amine or amide linkage. In one aspect of this preferred embodiment, the poly(ethylene glycol) is monomethoxy poly(ethylene glycol). In another aspect of this preferred embodiment, the poly(ethylene glycol) or poly(ethylene oxide) has a molecular weight between about 5 kDa and 30 kDa. Preferably, the poly(ethylene glycol) or poly(ethylene oxide) has a molecular weight between about 10 kDa and 20 kDa. Advantageously, the average number of strands of said poly(ethylene glycol) or poly(ethylene oxide) is between about 2 and 12 strands per uricase subunit. More advantageously, the average number of strands of said poly(ethylene glycol) or poly(ethylene oxide) is between about 6 and 10 per uricase subunit. Most advantageously, the average number of strands of said poly(ethylene glycol) or poly(ethylene oxide) is between about 7 and 9 per uricase subunit. Preferably, the poly(ethylene glycol) or poly(ethylene oxide) is linear. Alternatively, the poly(ethylene glycol) or poly(ethylene oxide) is branched.

[0012] The present invention also provides a pharmaceutical composition for lowering uric acid levels in a body fluid or tissue, comprising the uricase conjugate described above and a pharmaceutically acceptable carrier. Preferably, the composition is stabilized by lyophilization and dissolves upon reconstitution to provide solutions suitable for parenteral administration.

[0013] Another embodiment of the invention is a method for purifying uricase having reduced immunogenicity, comprising the step of separating uricase aggregates larger than octamers in uricase fractions, and excluding such aggregates from the purified uricase. Preferably, the separating step comprises the step of detecting aggregates larger than octamers from at least a portion of the uricase fractions and excluding the fractions containing the aggregates. Advantageously, the detecting step comprises measurement of light scattering.

[0014] The present invention also provides isolated uricase prepared by the method described above.

Brief Description of the Drawings**[0015]**

5 Figure 1 illustrates uricase activity, total protein and salt concentrations in fractions from a Pharmacia Biotech Mono Q (1 x 10 cm) anion exchange column. Uricase activity was measured at room temperature by monitoring the decrease in absorbance at 292 nm of 100 µM uric acid in 200 mM sodium borate, pH 9.2. Total protein was determined from the area under the curve of the absorbance peak of uricase in size-exclusion HPLC analyses.

10 Figure 2 illustrates size-exclusion HPLC analysis on a Pharmacia Superdex 200 column (1 x 30 cm) of the load and selected fractions from a preparative Mono Q chromatography of porcine uricase containing the mutations R291K and T301S (PKS uricase) showing data obtained by a light-scattering detector at 90°C (upper curves) and by absorbance at 276 nm (lower curves). The different signal strengths of the tetrameric, octameric and more highly aggregated forms of uricase in the unfractionated sample (load) and the various fractions are evident. The load was diluted 1/5 with Mono Q column buffer, fraction 5 was diluted 1/3 and fraction 6 was diluted 1/9. Fractions 5 and 6 were combined to form the "low salt pool."

15 Figure 3 illustrates size-exclusion analyses of fractions from the Mono Q column in Figure 1, showing data obtained by a light-scattering detector at 90° and by absorbance at 276 nm, as in Figure 2. The fractions shown in this figure were used to form the "high salt pool", from which PEG conjugates were prepared and injected into BALB/c mice. The resultant serum activities and immunologic responses in BALB/c mice are shown in Figures 5 and 6.

20 Figure 4 illustrates octamer content, determined by absorbance at 276 nm and by light scattering at 90°, calculated from the data in Figures 2 and 3, of unfractionated PKS uricase and of selected fractions from the preparative MonoQ column chromatography of PKS uricase (Figure 1).

25 Figure 5 illustrates UV assays, as in Figure 1, of uricase activity after a 4 hour incubation at 37°C, in sera drawn 24 hours after each of six weekly injections of 6 x 10-kDa PEG conjugates of PKS uricase or of pools from Mono Q column fractions.

30 Figure 6 illustrates ELISA analyses of IgG antibody formation against PEG conjugates of PKS uricase and against PEG conjugates of the pools of fractions from the Mono Q column shown in Figure 1, in sera drawn 24 hours after each of six weekly injections of female BALB/c mice with 0.2 mg of uricase protein per 20 grams of body weight. For each mouse, data from bleedings 24 hours after the first through sixth injections are shown from left to right. The assay conditions are described in Example 6. Data for the eight mice in each group were arranged in order of increasing immune response, from left to right.

Detailed Description of the Preferred Embodiments

35 **[0016]** Previous studies have shown that when a significant reduction in the immunogenicity and/ or antigenicity of uricase is achieved by conjugation with PEG (PEGylation), it is invariably associated with a substantial loss of uricolytic activity. The present invention includes the observation that traces of aggregates of urate oxidases larger than octamers substantially contribute to immunogenicity and the induction of accelerated clearance of PEG-uricase conjugates. This discovery is most likely applicable to proteins other than uricases, including interferons and growth factors.

40 **[0017]** The safety, convenience and cost- effectiveness of biopharmaceuticals are all adversely impacted by decreases in their potencies and the resultant need to increase the administered dose. Thus, there is a need for a safe and effective alternative means for lowering elevated levels of uric acid in body fluids, including blood and urine. The present invention provides a method for producing uricase that excludes uricase aggregates larger than octamers for use in the synthesis of PEG-uricase. This PEG-uricase retains all or nearly all of the uricolytic activity of the unmodified enzyme. The present invention also provides purified uricase substantially free of aggregates larger than octamers. The term "substantially free" indicates that the purified uricase comprises no more than about 2%, and preferably no more than about 1 % of aggregates larger than octamers.

45 **[0018]** The present invention provides a method for purifying uricase such that aggregates larger than octamers are excluded from the purified preparation. Because these larger aggregates are highly immunogenic, their presence in the purified uricase preparation is undesirable. The method involves monitoring column fractions by light scattering rather than or in addition to ultraviolet absorbance at 280 nm, because the aggregates may be too dilute to be detected by ultraviolet absorbance. The purified uricase is then conjugated to water- soluble polymers, preferably poly(ethylene glycols) or poly(ethylene oxides) as described in copending U.S. Application Serial No. 09/370,084.

50 **[0019]** The removal of aggregated uricase from a preparation consisting predominantly of tetrameric uricase can be accomplished by any of the methods known to those skilled in the art, including size-exclusion chromatography, ion-exchange chromatography, ultrafiltration through a microporous membrane and centrifugation, including ultracentrifugation. The separation method may include separation and analysis of fractions and the rejection or exclusion of those fractions containing excessive quantities of large aggregates. The resultant uricase preparation is better suited for the

synthesis of substantially non-immunogenic conjugates of uricase than is the unfractionated uricase. For chronic administration, it is important that PEG conjugates of proteins, e.g. PEG-uricase, have low immunogenicity and do not provoke progressively more rapid clearance from the bloodstream after repeated doses.

[0020] The invention also provides pharmaceutical compositions of the polymer-uricase conjugates. These conjugates are substantially non-immunogenic and retain at least 75%, preferably 85%, and more preferably 95% or more of the uricolytic activity of the unmodified enzyme. Uricases suitable for conjugation to water-soluble polymers include naturally occurring urate oxidases isolated from bacteria, fungi and the tissues of plants and animals, both vertebrates and invertebrates, as well as recombinant forms of uricase, including mutated, hybrid, and/or truncated enzymatically active variants of uricase. Water-soluble polymers suitable for use in the present invention include linear and branched poly(ethylene glycols) or poly(ethylene oxides), all commonly known as PEGs. Examples of branched PEG are the subject of U.S. Patent 5,643,575. One preferred example of linear PEG is monomethoxyPEG, of the general structure $\text{CH}_3\text{O} \cdot (\text{CH}_2\text{CH}_2\text{O})_n\text{H}$, where n varies from about 100 to about 2,300.

[0021] One embodiment of the present invention is a conjugate of urate oxidase (uricase) that retains at least about 75% of the uricolytic activity of unconjugated uricase and has substantially reduced immunogenicity. The uricase of this aspect of the invention may be recombinant. Whether recombinant or not, the uricase may be of mammalian origin. In one aspect of this embodiment, the uricase may be porcine, bovine or ovine liver uricase. In another aspect of this embodiment, the uricase may be chimeric. The chimeric uricase may contain portions of porcine liver and/or baboon liver uricase. For example, the chimeric uricase may be porcine uricase containing the mutations R291K and T301 S (PKS uricase). Alternatively, the uricase may be baboon liver uricase in which tyrosine 97 has been replaced by histidine, whereby the specific activity of the uricase may be increased by at least about 60%. The uricase of the invention, whatever the origin, may also be in a form that is truncated, either at the amino terminal, or at the carboxyl terminal, or at both terminals. Likewise, the uricase may be fungal or microbial uricase. In one aspect of this embodiment, the fungal or microbial uricase may be a naturally occurring or recombinant form of uricase from *Aspergillus flavus*, *Arthrobacter globiformis*, *Bacillus* sp. or *Candida utilis*. Alternatively, the uricase may be an invertebrate uricase, such as, for examples, a naturally occurring or recombinant form of uricase from *Drosophila melanogaster* or *Drosophila pseudoobscura*. The uricase of the invention may also be a plant uricase, for example, a naturally occurring or recombinant form of uricase from soybean root nodule (*Glycine max*). The PEG may have an average molecular weight between about 5 kDa and 100 kDa; preferably the PEG may have an average molecular weight between about 8 kDa and 60 kDa; more preferably, the PEG may have an average molecular weight between about 10 kDa and about 40 kDa, such as, for example, 10 to 20 kDa. The average number of covalently coupled strands of PEG may be 2 to 12 strands per uricase subunit; preferably, the average number of covalently coupled strands may be 6 to 10 per subunit; more preferably, the average number of strands of PEG may be 7 to 9 per subunit. In one aspect of this embodiment, the uricase may be tetrameric. The strands of PEG may be covalently linked to uricase via urethane (carbamate) linkages, secondary amine linkages, and/or amide linkages. When the uricase is a recombinant form of any of the uricases mentioned herein, the recombinant form may have substantially the sequence of the naturally occurring form.

[0022] One preferred mammalian uricase is recombinant pig-baboon chimeric uricase, composed of portions of the sequences of pig liver and baboon liver uricase, both of which were first determined by Wu, et al., (1989). One example of such a chimeric uricase contains the first 288 amino acids from the porcine sequence (SEQ ID NO: 1) and the last 16 amino acids from the baboon sequence (SEQ ID NO: 2). Hershfied, et al., International Publication WO 00/08196, 40 Urate Oxidase, published February 17, 2000. Since the latter sequence differs from the porcine sequence at only two positions, having a lysine (K) in place of arginine at residue 291 and a serine (S) in place of threonine at residue 301, this mutant is referred to as pig-K-S or PKS uricase (SEQ ID NO: 3). PKS uricase has one more lysine residue and, hence, one more potential site of PEGylation than either the porcine or baboon sequence. The sequences are shown in Table A.

[0023] The cDNAs for various mammalian uricases, including PKS uricase, were subcloned and the optimal conditions were determined for expression in *E. coli*, using standard methods. See Erlich, HA, (Ed.) (1989) PCR Technology. Principles and Applications for DNA Amplification. New York: Stockton Press; Sambrook, J, et al., (1989) Molecular Cloning. A Laboratory Manual, Second Edition. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press. The recombinant uricases were extracted, purified and their stability and activity were assessed using a modification of standard assays. See Fridovich, I, (1965) J Biol Chem 240:2491-2494; Nishimura, et al., (1979), and Examples 1 and 5.

[0024] In one embodiment of the invention, uricase may be conjugated via a biologically stable, nontoxic, covalent linkage to a relatively small number of strands of PEG. Such linkages may include urethane (carbamate) linkages, secondary amine linkages, and amide linkages. Various activated PEGs suitable for such conjugation are available commercially from Shearwater Polymers, Huntsville, AL.

[0025] For example, urethane linkages to uricase may be formed by incubating uricase in the presence of the succinimidyl carbonate (SC) or *p*-nitrophenyl carbonate (NPC) derivative of PEG. SC-PEG may be synthesized using the procedure described in U.S. Patent 5,612,460. NPC-PEG may be synthesized by reacting PEG with *p*-nitrophenyl chloroformate according to methods described in Veronese, FM, et al., (1985) Appl Biochem Biotechnol 11:141-152,

and in U.S. Patent 5,286,637. The methods described in the '637 patent are adapted to PEGs of higher molecular weight by adjusting the concentrations of the reactants to maintain similar stoichiometry. An alternative method of synthesis of NPC-PEG is described by Büttner, W., et al., East German Patent Specification DD 279 486 A1.

[0026] Amide linkages to uricase may be obtained using an N-hydroxysuccinimide ester of a carboxylic acid derivative of PEG (Shearwater Polymers). Secondary amine linkages may be formed using 2,2,2-trifluoroethanesulfonyl PEG (tresyl PEG; Shearwater Polymers) or by reductive alkylation using PEG aldehyde (Shearwater Polymers) and sodium cyanoborohydride.

[0027] In conjugates containing PEG with a molecular weight of 10 kDa, the maximum number of strands of PEG that were coupled per subunit, while retaining at least 75% of the uricolytic activity of the unmodified enzyme, was about 12 strands for mammalian uricases (e.g. PKS uricase, a mutein of porcine uricase; see assay conditions in Example 5). The latter extent of PEGylation corresponds to about 40% of the total amino groups. In one embodiment of the invention, the average number of strands of PEG coupled per uricase subunit is between about 2 and 12. In a preferred embodiment, the average number of strands of PEG coupled per uricase subunit is between about 6 and 10. In a more preferred embodiment, the average number of covalently linked strands of PEG per uricase subunit is between about 7 and 9. In another embodiment, the molecular weight of PEG used for the coupling reaction is between about 5 kDa and 30 kDa, preferably between about 10 kDa and 20 kDa.

[0028] There are several factors that may affect the choice of the optimal molecular weight and number of strands of PEG for coupling to a given form of uricase. In general, the reduction or elimination of immunogenicity without substantial loss of uricolytic activity may require the coupling of relatively more strands of PEG of lower molecular weight, compared to relatively fewer strands of PEG of higher molecular weight. Likewise, each different form of uricase may have a different optimum with respect to both the size and number of strands. The optimal number of strands of PEG and PEG molecular weight can be readily determined using the methods described herein.

[0029] When PEG conjugates of mammalian uricase were prepared from the purified tetrameric and octameric forms of the enzyme (containing four or eight subunit of approximately 35 kDa), they displayed profoundly reduced immunogenicity in mice, in contrast to the moderate immunogenicity of PEG conjugates of uricase preparations containing large aggregates (see Figure 6) and the very high immunogenicity of the unmodified enzyme.

[0030] Purified preparations of naturally occurring and recombinant uricases usually contain a mixture of very large aggregates of the enzyme, in addition to the tetrameric (140-kDa) and the octameric (280-kDa) forms. The percentage of each uricase preparation that is in either the tetrameric or octameric form generally varies from about 20 % to 95% (see Figures 2-4). Despite evidence that unPEGylated aggregates of several other proteins are highly immunogenic (see, e.g., Moore, WV, et al., (1980) J Clin Endocrinol Metab 51:691-697), previous studies of PEG-uricase do not describe any efforts to limit the content of aggregates, suggesting that the potential immunogenicity of the PEG-modified aggregates was not considered. On the basis of the observations of the present inventors, it appears likely that such aggregates were present in the enzyme preparations used for previous syntheses of PEG-uricase. Their presence may have rendered the task of preparing non-immunogenic conjugates more difficult. It also appears that the large losses of uricolytic activity observed in previous efforts to PEGylate uricase were related to the large number of strands of low molecular weight PEG that were coupled. On the other hand, the methods of uricase purification and PEGylation described herein permit the covalent attachment of as many as 12 strands of PEG per subunit while retaining more than 75% of the uricolytic activity, at least for certain uricases, e.g., PKS uricase (a mutein of porcine uricase) and the enzyme from thermophilic *Bacillus* sp.

[0031] In another preferred embodiment, substantially all large aggregates of the enzyme may be removed by ion-exchange chromatography (Figures 1-3) or size-exclusion chromatography at a pH between about 9 and 10.5, preferably 10.2, prior to conjugation of the resulting substantially aggregate-free preparation of uricase to PEG. The molecular weight of the uricase in each fraction from the preparative column may be monitored by any size-dependent analytical technique, including, for example, HPLC, conventional size-exclusion chromatography, centrifugation, light scattering, capillary electrophoresis or gel electrophoresis in a non-denaturing buffer. For aggregate-free uricase isolated using size-exclusion chromatography, fractions containing only the 140-kDa and 280-kDa forms of the enzyme may be pooled and used for conjugation to PEG. For tetrameric plus octameric uricase isolated using ion-exchange chromatography, fractions from the ion-exchange column may be analyzed with respect to size to determine which fractions contain substantial amounts of the tetrameric and octameric forms without the large aggregates detected by light scattering. In the purified product, the undesirable large aggregates may thus constitute as little as about 1%, or less, of the total uricase.

[0032] The results presented herein indicate that, even when extensively PEGylated, forms of PKS uricase larger than the octamer provoke accelerated clearance (Figure 5) and are somewhat immunogenic in mice (Figure 6). In contrast, conjugates prepared from uricase that is essentially free of large aggregates (detectable by light scattering) could be reinjected at least six times at one-week intervals with much less evidence of accelerated clearance rates (Figure 5) and without the detectable formation of antibodies, as measured by a sensitive enzyme-linked immunoassay (Figure 6). The use of highly purified tetrameric or octameric uricase further distinguishes the improved, conjugates of the present invention from the PEG-uricase preparations described previously. In contrast, the presence of a significant content of

large aggregates in the uricase preparations used by some previous investigators may have led them to couple large numbers of strands of PEG in efforts to suppress the immunogenicity. Consequently, the enzymatic activity of the resultant conjugates was decreased substantially.

[0033] The PEG-uricase conjugates of the present invention are useful for lowering the levels of uric acid in the body fluids and tissues of mammals, preferably humans, and can thus be used for treatment of elevated uric acid levels associated with conditions including gout, tophi, renal insufficiency, organ transplantation and malignant disease. PEG - uricase conjugates may be injected into a mammal having excessive uric acid levels by any of a number of routes, including intravenous, subcutaneous, intradermal, intramuscular and intraperitoneal routes. Alternatively, they may be aerosolized and inhaled. See Patton, JS, (1996) *Adv Drug Delivery Rev* 19:3-36 and U.S. Patent 5,458,135. The effective dose of PEG-uricase of the present invention will depend on the level of uric acid and the size of the individual. In one embodiment of this aspect of the invention, PEG-uricase is administered in a pharmaceutically acceptable excipient or diluent in an amount ranging from about 10 µg to about 1 g. In a preferred embodiment, the amount administered is between about 100 µg and 500 mg. More preferably, the conjugated uricase is administered in an amount between 1 mg and 100 mg, such as, for example, 5 mg, 20 mg or 50 mg. Masses given for dosage amounts of the embodiments refer to the amount of protein in the conjugate.

[0034] Pharmaceutical formulations containing PEG-uricase can be prepared by conventional techniques, e.g., as described in Gennaro, AR (Ed.) (1990) *Remington's Pharmaceutical Sciences*, 18th Edition, Easton, PA: Mack Publishing Co. Suitable excipients for the preparation of injectable solutions include, for example, phosphate buffered saline, lactated Ringer's solution, water, polyols and glycerol. Pharmaceutical compositions for parenteral injection comprise pharmaceutically acceptable sterile aqueous or non- aqueous liquids, dispersions, suspensions, or emulsions as well as sterile powders for reconstitution into sterile injectable solutions or dispersions just prior to use. These formulations may contain additional components, such as, for example, preservatives, solubilizers, stabilizers, wetting agents, emulsifiers, buffers, antioxidants and diluents.

[0035] PEG-uricase may also be provided as controlled-release compositions for implantation into an individual to continually control elevated uric acid levels in body fluids. For example, polylactic acid, polyglycolic acid, regenerated collagen, poly-L-lysine, sodium alginate, gellan gum, chitosan, agarose, multilamellar liposomes and many other conventional depot formulations comprise bioerodible or biodegradable materials that can be formulated with biologically active compositions. These materials, when implanted or injected, gradually break down and release the active material to the surrounding tissue. For example, one method of encapsulating PEG- uricase comprises the method disclosed in U.S. Patent 5,653,974. The use of bioerodible, biodegradable and other depot formulations is expressly contemplated in the present invention. The use of infusion pumps and matrix entrapment systems for delivery of PEG-uricase is also within the scope of the present invention. PEG-uricase may also advantageously be enclosed in micelles or liposomes. Liposome encapsulation technology is well known in the art. See, e.g., Lasic, D, et al., (Eds.) (1995) *Stealth Liposomes*. Boca Raton, FL: CRC Press.

[0036] The PEG-uricase pharmaceutical compositions of the invention will decrease the need for hemodialysis in patients at high risk of urate-induced renal failure, e.g., organ transplant recipients (see Venkataseshan, VS, et al., (1990) *Nephron* 56:317-321) and patients with some malignant diseases. In patients with large accumulations of crystalline urate (tophi), such pharmaceutical compositions will improve the quality of life more rapidly than currently available treatments.

[0037] The following examples, which are not to be construed as limiting the invention in any way, illustrate the various aspects disclosed above. These examples describe PEG-uricases prepared by coupling activated PEG (e.g., the *p*-nitrophenyl carbonate derivative) to a mutein of porcine uricases. These examples provide guidance to one of ordinary skill in the art for producing substantially non-immunogenic conjugates of uricase that retain at least about 75% of the uricolytic activity of the unmodified enzyme and are well suited for chronic administration.

Example 1

Preparative ion-exchange chromatography of uricase

[0038] Preparative ion-exchange chromatography was performed on a Fast Protein Liquid Chromatography (FPLC) apparatus (Amersham Pharmacia, Piscataway, NJ). The Mono Q column (1 x 10 cm, Amersham Pharmacia) was eluted with a gradient of 50 mM sodium carbonate, pH 10.3, 0.1 M NaCl (Buffer A) to 50 mM sodium carbonate, pH 10.3, 0.6 M NaCl (Buffer B) at a flow rate of 0.5 ml/min, except that the sample was loaded at a lower flow-rate. This technique was used to fractionate 25 mL of a solution of PKS uricase (pH 10.3). PKS uricase was obtained from Bio-Technology General Limited (Rehovot, Israel). The latter is recombinant porcine uricase in which one residue of lysine (K) and one residue of serine (S) have replaced one residue of arginine and one residue of threonine, respectively, in the parental porcine sequence (Lee et al. (1988) *Science* 239:1288-1291; Wu et al. (1989) *Proc. Natl. Acad. Sci. U. S. A.* 86: 9412-9416). After the sample was loaded, the column was washed with 100 mL of Buffer A. The peak of uricase began

to elute at the end of a 31-mL linear gradient of 0 to 26% Buffer B. Most of the uricase was eluted isocratically by 7 mL of buffer containing 26% Buffer B. The remainder of the recovered uricase was eluted by a linear 89-mL gradient of 26% to 100% buffer B. Fractions of 4 mL or 6 mL were collected. Aliquots of Fractions #4-11 were assayed for uricase and total protein (Figure 1) and were analyzed by size-exclusion high performance liquid chromatography (HPLC) as described in Example 2 (Figures 2 and 3). The remaining portions of Fractions #5-10 were coupled to PEG, as described in Example 3. Based on the results of the analyses in Example 2, the PEG conjugates of Fractions #5 and 6 were combined as the "Low-Salt Pool" and the PEG conjugates of Fractions #7-10 were combined as the "High-Salt Pool," as indicated in Figure 1.

10 Example 2

Size-exclusion chromatography of uricase monitored by light scattering and ultraviolet absorbance

[0039] Size-exclusion HPLC was performed at room temperature on a Superdex 200 column (1 x 30 cm, Amersham Pharmacia Biotech) on unfractionated PKS uricase and on selected fractions from the preparative Mono Q chromatography of PKS uricase at Example 1. The eluate from the absorbance monitor (UV 2000) of the Thermo Separations HPLC (Sunnyvale, CA) was analyzed by light scattering at 90° to the incident light, using a MiniDawn detector from Wyatt Technologies (Santa Barbara, CA).

[0040] The results shown in Figures 2-4 illustrate the resolution among the tetramer, octamer and larger aggregates of the uricase subunit and the different proportions of the signals detected from these forms of uricase in the various samples. Unlike the absorbance signal, which is directly proportional to the concentration, the light scattering signal is proportional to the product of the concentration times the size of the light scattering unit. The resultant sensitivity of the light scattering detector to very small amounts of highly aggregated uricase revealed the presence of the largest aggregates, which are eluted at or near the void volume (approximately 7 ml).

25 Example 3

Synthesis of PEG-uricase conjugates

[0041] Unfractionated PKS uricase (from Bio-Technology General Limited) and the uricase in fractions from the Mono Q column of Example 1 were coupled to 10-kDa PEG using the p-nitrophenyl carbonate derivative of PEG (NPC-PEG) obtained from Shearwater Polymers (Huntsville, AL). The preparation of NPC-PEG from PEG using phenylchloroformates has been described in several reports (e.g. Veronese, FM, et al., (1985) Appl Biochem Biotechnol 11:141-152; Kito, M, et al., (1996) J Clin Biochem Nutr 21:101-111) and NPC-PEG has been used for the synthesis of PEG-protein conjugates by previous investigators including the present inventors (e.g. Veronese *et al.*, *supra*; Sherman, MR, et al., in JM Harris, et al., (Eds.) Poly(ethylene glycol) Chemistry and Biological Applications. ACS Symposium Series 680 (pp. 155-176) Washington, DC: American Chemical Society). The number of strands of 10-kDa PEG coupled to each subunit of uricase was determined to be six by the method described by Kunitani, M, et al., (1991) J Chromatogr 588:125-137.

40 Example 4

In vivo serum persistence and immunogenicity of uricase and PEG-uricase

[0042] PEG conjugates of recombinant mammalian uricases, prepared according to the method of Example 3, were adjusted to 1 mg protein/mL in phosphate-buffered saline (PBS), pH 7.4, for injection. Samples were frozen and stored until analyzed or injected. Samples were warmed to 37°C for up to 1 hour prior to injection into groups of eight BALB/c female mice. The groups of mice had mean weights in the range of 18-22 g at the start of the studies.

[0043] The weights of all mice were monitored and evidence of adverse reactions to the injections or other evidence of ill health was recorded. Twenty-four hours after each of six weekly injections, the animals were anesthetized with ketamine and 100-200 µL of blood was obtained retro-orbitally, except at sacrifice (exsanguination), when a larger volume was collected. Serum was prepared from blood that had clotted for between 4 and 32 hours at 2-8°C. Sera were stored at -20°C. Sera were analyzed for uricolytic activity as described in Example 5 and analyzed for antibodies against uricases as described in Example 6.

Example 5Uricolytic activity assays of PEG-uricase in sera from mice injected with PEG-uricase

[0044] An activity assay based on ultraviolet light absorbance (UV assay) was performed with 100 μ M uric acid as the substrate in 200 mM sodium borate, pH 9.2, in a microplate adaptation of the method of I. Fridovich (J Biol Chem. (1965) 240:2491-2494). The decrease in absorbance at 292 nm was monitored for 15 minutes at room temperature in a 96-well plate with a UV-transparent bottom (Costar, Corning, NY), using a SpectraMAX 250 microplate reader from Molecular Devices (Sunnyvale, CA). The data were analyzed by finding the maximum slope (in milli-absorbance units per minute) of absorbance measurements made during the interval while between 10 and 40% of the substrate was oxidized. Results obtained with this assay are illustrated in Figures 1 and 5.

[0045] The mean half-life in sera of mice injected for the first time with PKS uricase coupled to six strands of 10-kDa PEG per subunit (6 x 10-kDa PEG PKS) was 29 \pm 4 hours, based on data from sera obtained 24 and 72 hours after the injection.

[0046] In separate experiments, it was established that the detectable uricolytic activity in the sera of mice injected with PEG-uricase declines during storage at -20°C and that maximal recovery of this activity is obtained by a 4-hour incubation at 37° prior to assay. Figure 5 shows that the recovery of uricolytic activity after repeated weekly injections of 6 x 10-kDa PEG PKS uricase was greatest when the enzyme was purified by Mono Q column chromatography, as in Example 1, prior to PEGylation according to the method of Example 3. Recovery was highest after the injection of conjugates prepared from the high-salt eluate pool of Example 1 (see Figure 1), which has the smallest content of the very large aggregates (see the light scattering profiles of Fractions 7-10 in Figure 3). Intermediate recovery was obtained with conjugates prepared from the low-salt eluate pool from the Mono Q column of Example 1, and the poorest recovery was obtained with conjugates made from unfractionated PKS uricase, which has the highest content of very large aggregates (see Figure 2). The same order of relative activities recovered in sera after repeated injections (high salt pool > low salt pool > unfractionated uricase) was observed regardless of whether the UV assay described above or a colorimetric assay adapted from P. Fossati et al. (J. Clin Chem (1980) 26:227-231), was used and regardless of whether the sera were incubated at 37°C before they were assayed.

Example 6Enzyme-linked immunosorbent assay (ELISA) of sera from mice injected with PEG-uricase

[0047] Non-competitive ELISA analyses were performed with porcine uricase bound to 96-well Immulon 2 plates (Dynex Technologies, from VWR Scientific, San Francisco, CA). The primary antisera were from mice injected with uricase or 6 x 10-kDa PEG conjugates prepared according to the method of Example 3. The secondary antibody was goat anti-mouse IgG coupled to horseradish peroxidase (Calbiochem-Novabiochem #401 253, La Jolla, CA) and the substrate was α -phenylenediamine dihydrochloride (Sigma P-9187, St. Louis. MO), as described by B. Porstmann et al. (J Clin. Chem. Clin. Biochem. (1981) 19:435-440).

[0048] Figure 6 illustrates the results of the non-competitive ELISA analyses. The results demonstrate that the 6 x 10-kDa PEG PKS uricase synthesized according to the method of Example 3 from the high-salt eluate from the Mono Q column of Example 1 (shown in Figure 1) did not produce detectable immune responses in any of the eight mice that received weekly injections for six weeks. A few mice injected with conjugates prepared from unfractionated PKS uricase according to the method of Example 3 showed low but detectable immune responses. The highest incidence of immune responses was in mice injected with conjugates prepared according to the method of Example 3 from the low-salt eluate pool from the Mono Q column of Example 1.

[0049] Without the benefit of the light scattering detector for the size-exclusion HPLC analyses, as described in Example 2, it would not have been apparent that the presence of the largest aggregates, not of the octameric form of uricase, is associated with progressively decreased recovery of PEG-uricase conjugates after repeated injections, as observed in Example 5 (Figure 5) and with an increase in immunogenicity in BALB/c mice, as observed in Example 6 (Figure 6).

These results have important implications for the specifications of the uricase used as a starting material for the production of PEG-uricase for clinical use.

TABLE A: SEQUENCES

SEQ ID NO: 1

5 Met Ala His Tyr Arg Asn Asp Tyr Lys Lys Asn Asp Glu Val Glu Phe
 1 5 10 15

10 Val Arg Thr Gly Tyr Gly Lys Asp Met Ile Lys Val Leu His Ile Gln
 20 25 30

15 Arg Asp Gly Lys Tyr His Ser Ile Lys Glu Val Ala Thr Ser Val Gln
 35 40 45

20 Leu Thr Leu Ser Ser Lys Lys Asp Tyr Leu His Gly Asp Asn Ser Asp
 50 55 60

25 Val Ile Pro Thr Asp Thr Ile Lys Asn Thr Val Asn Val Leu Ala Lys
 65 70 75 80

30 Phe Lys Gly Ile Lys Ser Ile Glu Thr Phe Ala Val Thr Ile Cys Glu
 85 90 95

35 His Phe Leu Ser Ser Phe Lys His Val Ile Arg Ala Gln Val Tyr Val
 100 105 110

40 Glu Glu Val Pro Trp Lys Arg Phe Glu Lys Asn Gly Val Lys His Val
 115 120 125

45 His Ala Phe Ile Tyr Thr Pro Thr Gly Thr His Phe Cys Glu Val Glu
 130 135 140

50 Gln Ile Arg Asn Gly Pro Pro Val Ile His Ser Gly Ile Lys Asp Leu
 145 150 155 160

55 Lys Val Leu Lys Thr Thr Gln Ser Gly Phe Glu Gly Phe Ile Lys Asp
 165 170 175

60 Gln Phe Thr Thr Leu Pro Glu Val Lys Asp Arg Cys Phe Ala Thr Gln
 180 185 190

65 Val Tyr Cys Lys Trp Arg Tyr His Gln Gly Arg Asp Val Asp Phe Glu
 195 200 205

70 Ala Thr Trp Asp Thr Val Arg Ser Ile Val Leu Gln Lys Phe Ala Gly
 210 215 220

EP 1 254 237 B1

5 Pro Tyr Asp Lys Gly Glu Tyr Ser Pro Ser Val Gln Lys Thr Leu Tyr
225 230 235 240

Asp Ile Gln Val Leu Thr Leu Gly Gln Val Pro Glu Ile Glu Asp Met
245 250 255

10 Glu Ile Ser Leu Pro Asn Ile His Tyr Leu Asn Ile Asp Met Ser Lys
260 265 270

15 Met Gly Leu Ile Asn Lys Glu Glu Val Leu Leu Pro Leu Asp Asn Pro
275 280 285

20 Tyr Gly Arg Ile Thr Gly Thr Val Lys Arg Lys Leu Thr Ser Arg Leu
290 295 300

SEQ ID NO: 2

25 Met Ala Asp Tyr His Asn Asn Tyr Lys Lys Asn Asp Glu Leu Glu Phe
1 5 10 15

30 Val Arg Thr Gly Tyr Gly Lys Asp Met Val Lys Val Leu His Ile Gln
20 25 30

Arg Asp Gly Lys Tyr His Ser Ile Lys Glu Val Ala Thr Ser Val Gln
35 40 45

35 Leu Thr Leu Ser Ser Lys Lys Asp Tyr Leu His Gly Asp Asn Ser Asp
50 55 60

40 Ile Ile Pro Thr Asp Thr Ile Lys Asn Thr Val His Val Leu Ala Lys
65 70 75 80

45 Phe Lys Gly Ile Lys Ser Ile Glu Ala Phe Gly Val Asn Ile Cys Glu
85 90 95

Tyr Phe Leu Ser Ser Phe Asn His Val Ile Arg Ala Gln Val Tyr Val
100 105 110

50 Glu Glu Ile Pro Trp Lys Arg Leu Glu Lys Asn Gly Val Lys His Val
115 120 125

55 His Ala Phe Ile His Thr Pro Thr Gly Thr His Phe Leu Glu Val Glu
130 135 140

• Gln Leu Arg Ser Gly Pro Pro Val Ile His Ser Gly Ile Lys Asp Leu
 145 150 155 160

5

Lys Val Leu Lys Thr Thr Gln Ser Gly Phe Glu Gly Phe Ile Lys Asp
 165 170 175

10 Gln Phe Thr Thr Lys Pro Glu Val Lys Asp Arg Cys Phe Ala Thr Gln
 180 185 190

15 Val Tyr Cys Lys Trp Arg Tyr His Gln Cys Arg Asp Val Asp Phe Glu
 195 200 205

20 Ala Thr Trp Gly Thr Ile Arg Asp Leu Val Leu Glu Lys Phe Ala Gly
 210 215 220

25 Pro Tyr Asp Lys Gly Glu Tyr Ser Pro Ser Val Gln Lys Thr Leu Tyr
 225 230 235 240

30 Asp Ile Gln Val Leu Ser Leu Ser Arg Val Pro Glu Ile Glu Asp Met
 245 250 255

35 Glu Ile Ser Leu Pro Asn Ile His Tyr Phe Asn Ile Asp Met Ser Lys
 260 265 270

Met Gly Leu Ile Asn Lys Glu Glu Val Leu Leu Pro Leu Asp Asn Pro
 275 280 285

35

Tyr Gly Lys Ile Thr Gly Thr Val Lys Arg Lys Leu Ser Ser Arg Leu
 290 295 300

40

SEQ ID NO:3

45 Met Ala His Tyr Arg Asn Asp Tyr Lys Lys Asn Asp Glu Val Glu Phe
 1 5 10 15

50 Val Arg Thr Gly Tyr Gly Lys Asp Met Ile Lys Val Leu His Ile Gln
 20 25 30

Arg Asp Gly Lys Tyr His Ser Ile Lys Glu Val Ala Thr Ser Val Gln
 35 40 45

55 Leu Thr Leu Ser Ser Lys Lys Asp Tyr Leu His Gly Asp Asn Ser Asp
 50 55 60

EP 1 254 237 B1

Val Ile Pro Thr Asp Thr Ile Lys Asn Thr Val Asn Val Leu Ala Lys
65 . 70 75 80

5 Phe Lys Gly Ile Lys Ser Ile Glu Thr Phe Ala Val Thr Ile Cys Glu
85 90 95

10 His Phe Leu Ser Ser Phe Lys His Val Ile Arg Ala Gln Val Tyr Val
100 105 110

15 Glu Glu Val Pro Trp Lys Arg Phe Glu Lys Asn Gly Val Lys His Val
115 120 125

His Ala Phe Ile Tyr Thr Pro Thr Gly Thr His Phe Cys Glu Val Glu
130 135 140

20 Gln Ile Arg Asn Gly Pro Pro Val Ile His Ser Gly Ile Lys Asp Leu
145 150 155 160

25 Lys Val Leu Lys Thr Thr Gln Ser Gly Phe Glu Gly Phe Ile Lys Asp
165 170 175

30 Gln Phe Thr Thr Leu Pro Glu Val Lys Asp Arg Cys Phe Ala Thr Gln
180 185 190

35 Val Tyr Cys Lys Trp Arg Tyr His Gln Gly Arg Asp Val Asp Phe Glu
195 200 205

40 Ala Thr Trp Asp Thr Val Arg Ser Ile Val Leu Gln Lys Phe Ala Gly
210 215 220

45 Pro Tyr Asp Lys Gly Glu Tyr Ser Pro Ser Val Gln Lys Thr Leu Tyr
225 230 235 240

Asp Ile Gln Val Leu Thr Leu Gly Gln Val Pro Glu Ile Glu Asp Met
245 250 255

50 Glu Ile Ser Leu Pro Asn Ile His Tyr Leu Asn Ile Asp Met Ser Lys
260 265 270

Met Gly Leu Ile Asn Lys Glu Glu Val Leu Leu Pro Leu Asp Asn Pro
275 280 285

55 Tyr Gly Lys Ile Thr Gly Thr Val Lys Arg Lys Leu Ser Ser Arg Leu
290 295 300

Claims

1. A uricase conjugate comprising a purified urate oxidase (uricase) conjugated to poly(ethylene glycol) or poly(ethylene oxide), wherein said uricase contains purified tetrameric and octameric forms of the uricase preparable by removing aggregates larger than octamers, wherein said uricase contains no more than about 2% of aggregates larger than octamers.
2. The uricase conjugate of Claim 1, wherein the uricase is mammalian uricase.
3. The uricase conjugate of Claim 2, wherein the uricase is porcine liver, bovine liver or ovine liver uricase.
4. The uricase conjugate of Claim 1, wherein the uricase is recombinant.
5. The uricase conjugate of Claim 4, wherein the uricase has substantially the sequence of porcine, bovine, ovine or baboon liver uricase.
6. The uricase conjugate of Claim 1, wherein the uricase is chimeric.
7. The uricase conjugate of Claim 6, wherein the chimeric uricase contains portions of porcine liver and baboon liver uricase.
8. The uricase conjugate of Claim 6, wherein the chimeric uricase is porcine uricase in which arginine residue 291 of SEQ ID NO:1 has been replaced by lysine (R291 K) and threonine residue 301 of SEQ ID NO:1 has been replaced by serine (T301 S) (PKS uricase).
9. The uricase conjugate of Claim 4, wherein the uricase has the sequence of baboon liver uricase having SEQ ID NO:2 in which tyrosine residue 97 of SEQ ID NO:2 has been replaced by histidine.
10. The uricase conjugate of Claim 4, wherein the uricase comprises an amino terminus and a carboxyl terminus, and wherein the uricase is truncated at one terminus or both termini.
11. The uricase conjugate of Claim 1, wherein the uricase is a fungal or microbial uricase.
12. The uricase conjugate of Claim 11, wherein the fungal or microbial uricase is isolated from *Aspergillus flavus*, *Arthrobacter globiformis*, *Bacillus sp.* or *Candida utilis*, or is a recombinant enzyme having substantially the sequence of one of said uricases.
13. The uricase conjugate of Claim 1, wherein the uricase is an invertebrate uricase.
14. The uricase conjugate of Claim 13, wherein the invertebrate uricase is isolated from *Drosophila melanogaster* or *Drosophila pseudoobscura*, or is a recombinant enzyme having substantially the sequence of one of said uricases.
15. The uricase conjugate of Claim 1, wherein the uricase is a plant uricase.
16. The uricase conjugate of Claim 15, wherein the plant uricase is isolated from root nodules of Glycine max or is a recombinant enzyme having substantially the sequence of said uricase.
17. The uricase conjugate of Claim 1, wherein said poly(ethylene glycol) is monomethoxy poly(ethylene glycol).
18. The uricase conjugate of Claim 1, wherein said uricase is conjugated to said poly(ethylene glycol) or poly(ethylene oxide) via a linkage selected from the group consisting of urethane (carbamate), secondary amine and amide.
19. The uricase conjugate of Claim 1, wherein said poly(ethylene glycol) or poly(ethylene oxide) has a molecular weight between about 5 kDa and 30 kDa.
20. The uricase conjugate of Claim 19, wherein said poly(ethylene glycol) or poly(ethylene oxide) has a molecular weight between about 10 kDa and 20 kDa.

21. The uricase conjugate of Claim 1, wherein the average number of strands of said poly(ethylene glycol) or poly(ethylene oxide) is between about 2 and 12 per uricase subunit.
- 5 22. The uricase conjugate of Claim 21, wherein the average number of strands of said poly(ethylene glycol) or poly(ethylene oxide) is between about 6 and 10 per uricase subunit.
23. The uricase conjugate of Claim 22, wherein the average number of strands of poly(ethylene glycol) or poly(ethylene oxide) is between about 7 and 9 per uricase subunit.
- 10 24. The uricase conjugate of Claim 1, wherein the poly(ethylene glycol) or poly(ethylene oxide) is linear.
25. The uricase conjugate of Claim 1, wherein the poly(ethylene glycol) or poly(ethylene oxide) is branched.
- 15 26. A pharmaceutical composition for lowering uric acid levels in a body fluid or tissue, comprising the conjugate of Claim 1 and a pharmaceutically acceptable carrier.
27. The pharmaceutical composition of Claim 26, wherein said composition is stabilized by lyophilization and dissolves upon reconstitution to provide solutions suitable for parenteral administration.
- 20 28. A method for purifying uricase having reduced immunogenicity, comprising the step of separating uricase aggregates larger than octamers in uricase fractions, and excluding such aggregates from the purified uricase, wherein the purified uricase contains purified tetrameric and octameric forms of the uricase preparable by removing aggregates larger than octamers, wherein said uricase contains no more than about 2% of aggregates larger than octamers.
- 25 29. The method of Claim 28, wherein said separating step is selected from the group consisting of ion-exchange chromatography, size-exclusion chromatography, and ultrafiltration.
- 30 30. The method of Claim 28, wherein said separating step comprises the step of detecting aggregates larger than octamers in uricase fractions and excluding said fractions containing said aggregates.
31. Isolated uricase prepared by the method of Claim 28.
- 35 32. A method for making a uricase conjugate, said method comprising the steps of:
applying a solution of uricase comprising tetrameric uricase, octameric uricase, and uricase aggregates larger than octamers to at least one separation column;
recovering from said column one or more fractions that contain isolated tetrameric and octameric uricase preparable by removing aggregates larger than octamers, wherein said uricase contains no more than about 2% of aggregates larger than octamers; and
40 PEGylating said uricase.
33. The method of Claim 32, wherein said separation column is selected from the group consisting of ion exchange column and size exclusion column.
- 45 34. The method of Claim 33, wherein said solution of said uricase is applied to said column at a pH of 10.2.
35. The method of Claim 33, further comprising the step of analyzing said fractions to determine at least one property selected from the group consisting of presence of said octameric uricase and absence of uricase aggregates that are larger than octamers.
- 50 36. The method of Claim 35, wherein said analyzing step comprises at least one analysis technique selected from the group consisting of chromatography, centrifugation, light scattering and electrophoresis.
37. The method of Claim 36, wherein said chromatography is high performance liquid chromatography.
- 55 38. A method for purifying uricase having reduced immunogenicity, comprising: separating uricase aggregates larger than octamers into uricase fractions; and recovering a fraction, wherein said fraction contains octameric uricase but does not contain uricase aggregates larger than octamers.

39. The uricase conjugate of Claim 1, wherein at least 2.5% of said uricase is in the octameric form.

Patentansprüche

- 5 1. Urikasekonjugat, umfassend eine gereinigte Uratoxidase (Urikase), konjugiert an Poly(ethylenglykol) oder Poly(ethylenoxid), wobei die Urikase gereinigte tetramere und oktamere Formen der Urikase, welche durch Entfernen von Aggregaten, die größer als Oktamere sind, hergestellt werden können, enthält, wobei die Urikase nicht mehr als etwa 2 % Aggregate, die größer als Oktamere sind, enthält.
- 10 2. Urikasekonjugat nach Anspruch 1, wobei die Urikase Säugetierurikase ist.
- 15 3. Urikasekonjugat nach Anspruch 2, wobei die Urikase Schweineleber-, Rinderleber- oder Schafleberurikase ist.
- 20 4. Urikasekonjugat nach Anspruch 1, wobei die Urikase rekombinant ist.
- 25 5. Urikasekonjugat nach Anspruch 4, wobei die Urikase im Wesentlichen die Sequenz von Schweine-, Rinder-, Schaf- oder Pavianleberurikase aufweist.
- 30 6. Urikasekonjugat nach Anspruch 1, wobei die Urikase chimär ist.
- 35 7. Urikasekonjugat nach Anspruch 6, wobei die chimäre Urikase Teile von Schweineleber- und Pavianleberurikase enthält.
- 40 8. Urikasekonjugat nach Anspruch 6, wobei die chimäre Urikase Schweineurikase ist, in welcher Argininrest 291 von SEQ ID Nr.: 1 durch Lysin (R291 K) ersetzt wurde und Threoninrest 301 von SEQ ID Nr.: 1 durch Serin (T301 S) ersetzt wurde (PKS-Urikase).
- 45 9. Urikasekonjugat nach Anspruch 4, wobei die Urikase die Sequenz von Pavianleberurikase mit SEQ ID Nr.: 2, in welcher Tyrosinrest 97 von SEQ ID Nr.: 2 durch Histidin ersetzt wurde, aufweist.
- 50 10. Urikasekonjugat nach Anspruch 4, wobei die Urikase einen Aminoterminus und einen Carboxylterminus umfasst und wobei die Urikase an einem Terminus oder beiden Terminen trunkiert ist.
- 55 11. Urikasekonjugat nach Anspruch 1, wobei die Urikase eine Pilz- oder Mikrobenurikase ist.
- 60 12. Urikasekonjugat nach Anspruch 11, wobei die Pilz- oder Mikrobenurikase aus *Aspergillus flavus*, *Arthrobacter globiformis*, *Bacillus sp.* oder *Candida utilis* isoliert ist oder ein rekombinantes Enzym mit im Wesentlichen der Sequenz von einer der Urikassen ist.
- 65 13. Urikasekonjugat nach Anspruch 1, wobei die Urikase eine Invertebraturikase ist.
- 70 14. Urikasekonjugat nach Anspruch 13, wobei die Invertebraturikase aus *Drosophila melanogaster* oder *Drosophila pseudoobscura* isoliert ist oder ein rekombinantes Enzym mit im Wesentlichen der Sequenz von einer der Urikassen ist.
- 75 15. Urikasekonjugat nach Anspruch 1, wobei die Urikase eine Pflanzenurikase ist.
- 80 16. Urikasekonjugat nach Anspruch 15, wobei die Pflanzenurikase aus Wurzelknollen von *Glycine max* isoliert ist oder ein rekombinantes Enzym mit im Wesentlichen der Sequenz der Urikase ist.
- 85 17. Urikasekonjugat nach Anspruch 1, wobei das Poly(ethylenglykol) Monomethoxypoly(ethylenglykol) ist.
- 90 18. Urikasekonjugat nach Anspruch 1, wobei die Urikase an das Poly(ethylenglykol) oder Poly(ethylenoxid) über eine Bindung, ausgewählt aus der Gruppe, welche aus Urethan(carbamat), sekundärem Amin und Amid besteht, konjugiert ist.
- 95 19. Urikasekonjugat nach Anspruch 1, wobei das Poly(ethylenglykol) oder Poly(ethylenoxid) ein Molekulargewicht von

zwischen etwa 5 kDa und 30 kDa aufweist.

20. Urikasekonjugat nach Anspruch 19, wobei das Poly(ethylenglykol) oder Poly(ethylenoxid) ein Molekulargewicht von zwischen etwa 10 kDa und 20 kDa aufweist.

5 21. Urikasekonjugat nach Anspruch 1, wobei die mittlere Anzahl von Strängen des Poly(ethylenglykol)s oder Poly(ethylenoxid)s zwischen etwa 2 und 12 pro Urikaseuntereinheit liegt.

10 22. Urikasekonjugat nach Anspruch 21, wobei die mittlere Anzahl von Strängen des Poly(ethylenglykol)s oder Poly(ethylenoxid)s zwischen etwa 6 und 10 pro Urikaseuntereinheit liegt.

- 15 23. Urikasekonjugat nach Anspruch 22, wobei die mittlere Anzahl von Strängen von Poly(ethylenglykol) oder Poly(ethylenoxid) zwischen etwa 7 und 9 pro Urikaseuntereinheit liegt.

20 24. Urikasekonjugat nach Anspruch 1, wobei das Poly(ethylenglykol) oder Poly(ethylenoxid) linear ist.

- 25 25. Urikasekonjugat nach Anspruch 1, wobei das Poly(ethylenglykol) oder Poly(ethylenoxid) verzweigt ist.

20 26. Arzneimittel zur Erniedrigung von Harnsäurelevels in einem Körperfluid oder Gewebe, umfassend das Konjugat nach Anspruch 1 und einen pharmazeutisch verträglichen Träger.

- 25 27. Arzneimittel nach Anspruch 26, wobei die Zusammensetzung durch Lyophilisation stabilisiert ist und sich bei Rekonstitution löst, wodurch Lösungen entstehen, welche für parenterale Verabreichung geeignet sind.

30 28. Verfahren zur Reinigung von Urikase mit verringriger Immunogenität, umfassend den Schritt von Abtrennen von Urikaseaggregaten, welche größer als Oktamere sind, in Urikasefraktionen und Ausschließen von solchen Aggregaten aus der gereinigten Urikase, wobei die gereinigte Urikase gereinigte tetramere und oktamere Formen der Urikase enthält, welche durch Entfernen von Aggregaten, die größer als Oktamere sind, hergestellt werden können, wobei die Urikase nicht mehr als etwa 2 % Aggregate, die größer als Oktamere sind, enthält.

35 29. Verfahren nach Anspruch 28, wobei der Abtrennungsschritt aus der Gruppe ausgewählt ist, welche aus Ionenaustauschchromatographie, Größenausschlusschromatographie und Ultrafiltration besteht.

35 30. Verfahren nach Anspruch 28, wobei der Abtrennungsschritt den Schritt von Nachweisen von Aggregaten, welche größer als Oktamere sind, in Urikasefraktionen und Ausschließen der Fraktionen, welche die Aggregate enthalten, umfasst.

- 40 31. Isolierte Urikase, hergestellt durch das Verfahren nach Anspruch 28.

40 32. Verfahren zur Herstellung eines Urikasekonjugats, wobei das Verfahren die folgenden Schritte umfasst:

45 das Aufbringen einer Lösung von Urikase, umfassend tetramere Urikase, oktamere Urikase und Urikaseaggregate, welche größer als Oktamere sind, auf mindestens eine Trennsäule;

45 das Gewinnen einer oder mehrerer Fraktionen von der Säule, welche isolierte tetramere und oktamere Urikase enthalten, die durch Entfernen von Aggregaten, die größer als Oktamere sind, hergestellt werden kann, wobei die Urikase nicht mehr als etwa 2 % Aggregate, welche größer als Oktamere sind, enthält; und das PEGylieren der Urikase.

50 33. Verfahren nach Anspruch 32, wobei die Trennsäule aus der Gruppe ausgewählt ist, welche aus Ionenaustauschsäule und Größenausschlusssäule besteht.

- 55 34. Verfahren nach Anspruch 33, wobei die Lösung der Urikase auf die Säule bei einem pH-Wert von 10,2 aufgebracht wird.

55 35. Verfahren nach Anspruch 33, ferner umfassend den Schritt von Analysieren der Fraktionen, um mindestens eine Eigenschaft zu bestimmen, welche aus der Gruppe ausgewählt ist, die aus der Gegenwart der oktameren Urikase und der Abwesenheit von Urikaseaggregaten, welche größer als Oktamere sind, besteht.

36. Verfahren nach Anspruch 35, wobei der Analyseschritt mindestens eine Analysetechnik umfasst, welche aus der Gruppe ausgewählt ist, die aus Chromatographie, Zentrifugation, Lichtstreuung und Elektrophorese besteht.
- 5 37. Verfahren nach Anspruch 36, wobei die Chromatographie Hochleistungsflüssigkeitschromatographie ist.
38. Verfahren zum Reinigen von Urikase mit verringelter Immunogenität, umfassend: das Trennen von Urikaseaggregaten, welche größer als Oktamere sind, in Urikasefraktionen; und das Gewinnen einer Fraktion, wobei die Fraktion oktamere Urikase enthält, aber keine Urikaseaggregate, welche größer als Oktamere sind, enthält.
- 10 39. Urikasekonjugat nach Anspruch 1, wobei mindestens 2,5 % der Urikase in der oktameren Form vorliegt.

Revendications

- 15 1. Conjugué d'uricase comprenant une urate oxydase (uricase) purifiée conjuguée à du polyéthylèneglycol ou à du poly(oxycde d'éthylène), dans lequel ladite uricase contient des formes tétramères et octamères purifiées de l'uricase qui peuvent être préparées par élimination des agrégats plus gros que des octamères, dans lequel ladite uricase ne contient pas plus d'environ 2 % d'agrégats plus gros que des octamères.
- 20 2. Conjugué d'uricase selon la revendication 1, dans lequel l'uricase est une uricase de mammifère.
3. Conjugué d'uricase selon la revendication 2, dans lequel l'uricase est une uricase de foie de porc, de foie de bovin ou de foie d'ovin.
- 25 4. Conjugué d'uricase selon la revendication 1, dans lequel l'uricase est recombinée.
5. Conjugué d'uricase selon la revendication 4, dans lequel l'uricase a pratiquement la séquence d'uricase de foie de porc, de bovin, d'ovin ou de babouin.
- 30 6. Conjugué d'uricase selon la revendication 1, dans lequel l'uricase est chimère.
7. Conjugué d'uricase selon la revendication 6, dans lequel l'uricase chimère contient des portions d'uricase de foie de porc et de foie de babouin.
- 35 8. Conjugué d'uricase selon la revendication 6, dans lequel l'uricase chimère est une uricase de porc dans laquelle le résidu d'arginine 291 de la SEQ ID NO : 1 a été remplacé par une lysine (R291 K) et le résidu de thréonine 301 de la SEQ ID NO : 1 a été remplacé par une sérine (T301 S) (uricase PKS).
- 40 9. Conjugué d'uricase selon la revendication 4, dans lequel l'uricase a la séquence de l'uricase de foie de babouin ayant la SEQ ID NO : 2 dans laquelle le résidu de tyrosine 97 de la SEQ ID NO : 2 a été remplacé par une histidine.
10. Conjugué d'uricase selon la revendication 4, dans lequel l'uricase comprend une extrémité amino et une extrémité carboxyle, et dans lequel l'uricase est tronquée à une extrémité ou aux deux extrémités.
- 45 11. Conjugué d'uricase selon la revendication 1, dans lequel l'uricase est une uricase fongique ou microbienne.
12. Conjugué d'uricase selon la revendication 11, dans lequel l'uricase fongique ou microbienne est isolée à partir d'*Aspergillus flavus*, *Arthrobacter globiformis*, *Bacillus sp.* ou *Candida utilis*, ou est une enzyme recombinée ayant pratiquement la séquence de l'une desdites uricases.
- 50 13. Conjugué d'uricase selon la revendication 1, dans lequel l'uricase est une uricase d'invertébré.
14. Conjugué d'uricase selon la revendication 13, dans laquelle l'uricase d'invertébré est isolée à partir de *Drosophila melanogaster* ou *Drosophila pseudoobscura*, ou est une enzyme recombinée ayant pratiquement la séquence de l'une desdites uricases.
- 55 15. Conjugué d'uricase selon la revendication 1, dans lequel l'uricase est une uricase végétale.

16. Conjugué d'uricase selon la revendication 15, dans lequel l'uricase végétale est isolée à partir de nodules de racines de Glycine max ou est une enzyme recombinée ayant pratiquement la séquence de ladite uricase.
- 5 17. Conjugué d'uricase selon la revendication 1, dans lequel ledit polyéthylèneglycol est du polyéthylèneglycol mono-méthoxylé.
- 10 18. Conjugué d'uricase selon la revendication dans lequel ladite uricase est conjuguée audit polyéthylèneglycol ou poly(oxyde d'éthylène) par une liaison choisie dans le groupe constitué par un uréthane (carbamate), une amine secondaire et un amide.
- 15 19. Conjugué d'uricase selon la revendication 1, dans lequel ledit polyéthylèneglycol ou poly(oxyde d'éthylène) a une masse moléculaire comprise entre environ 5 kDa et 30 kDa.
- 20 20. Conjugué d'uricase selon la revendication 19, dans lequel ledit polyéthylèneglycol ou poly(oxyde d'éthylène) a une masse moléculaire comprise entre environ 10 kDa et 20 kDa.
- 25 21. Conjugué d'uricase selon la revendication 1, dans lequel le nombre moyen de brins dudit polyéthylèneglycol ou poly(oxyde d'éthylène) est compris entre environ 2 et 12 par sous-unité d'uricase.
22. Conjugué d'uricase selon la revendication 21, dans lequel le nombre moyen de brins dudit polyéthylèneglycol ou poly(oxyde d'éthylène) est compris entre environ 6 et 10 par sous-unité d'uricase.
23. Conjugué d'uricase selon la revendication 22, dans lequel le nombre moyen de brins de polyéthylèneglycol ou poly(oxyde d'éthylène) est compris entre environ 7 et 9 par sous-unité d'uricase.
24. Conjugué d'uricase selon la revendication 1, dans lequel le polyéthylèneglycol ou poly(oxyde d'éthylène) est linéaire.
- 25 25. Conjugué d'uricase selon la revendication 1, dans lequel le polyéthylèneglycol ou poly(oxyde d'éthylène) est ramifié.
- 30 26. Composition pharmaceutique pour abaisser les taux d'acide urique dans un fluide corporel ou tissu corporel, comprenant le conjugué de la revendication 1 et un véhicule acceptable en pharmacie.
27. Composition pharmaceutique selon la revendication 26, laquelle composition est stabilisée par lyophilisation et se dissout lors d'une reconstitution pour former des solutions adaptées à une administration par voie parentérale.
- 35 28. Procédé pour purifier une uricase ayant une immunogénicité réduite, comprenant l'étape consistant à séparer les agrégats d'uricase plus gros que des octamères dans des fractions d'uricase, et à exclure ces agrégats de l'uricase purifiée, dans lequel l'uricase purifiée contient des formes tétramères et octamères purifiées de l'uricase qui peuvent être préparées par élimination des agrégats plus gros que des octamères, dans lequel ladite uricase ne contient pas plus d'environ 2 % d'agrégats plus gros que des octamères.
- 40 29. Procédé selon la revendication 28, dans lequel ladite étape de séparation est choisie dans le groupe constitué par une chromatographie par échange d'ions, une chromatographie par exclusion de tailles et une ultrafiltration.
- 45 30. Procédé selon la revendication 28, dans lequel ladite étape de séparation comprend l'étape consistant à détecter les agrégats plus gros que des octamères dans les fractions d'uricase et à exclure lesdites fractions contenant lesdits agrégats.
31. Uricase isolée préparée par le procédé de la revendication 28.
- 50 32. Procédé pour produire un conjugué d'uricase, ledit procédé comprenant les étapes consistant à :
- appliquer une solution d'uricase comprenant de l'uricase tétramère, de l'uricase octamère et des agrégats d'uricase plus gros que des octamères, à au moins une colonne de séparation ;
- 55 récupérer à partir de ladite colonne une ou plusieurs fractions qui contiennent de l'uricase tétramère et octamère isolée pouvant être préparée par élimination des agrégats plus gros que des octamères, ladite uricase ne contenant pas plus d'environ 2 % d'agrégats plus gros que des octamères ; et soumettre ladite uricase à une PEGylation.

33. Procédé selon la revendication 32, dans lequel ladite colonne de séparation est choisie dans le groupe constitué par une colonne échangeuse d'ions et une colonne d'exclusion de tailles.

5 34. Procédé selon la revendication 33, dans lequel ladite solution de ladite uricase est appliquée à ladite colonne à un pH de 10,2.

10 35. Procédé selon la revendication 33, comprenant en outre l'étape consistant à analyser lesdites fractions pour déterminer au moins une propriété choisie dans le groupe constitué par la présence de ladite uricase octamère et l'absence d'agrégats d'uricase plus gros que des octamères.

15 36. Procédé selon la revendication 35, dans lequel ladite étape d'analyse comprend au moins une technique d'analyse choisie dans le groupe constitué par la chromatographie, la centrifugation, la diffusion de lumière et l'électrophorèse.

20 37. Procédé selon la revendication 36, dans lequel ladite chromatographie est une chromatographie liquide à haute performance.

25 38. Procédé pour purifier une uricase ayant une immunogénicité réduite, consistant à : séparer les agrégats d'uricase plus gros que des octamères dans les fractions d'uricase ; et récupérer une fraction, dans lequel ladite fraction contient de l'uricase octamère, mais ne contient pas d'agrégats d'uricase plus gros que des octamères.

30 39. Conjugué d'uricase selon la revendication 1, dans lequel au moins 2,5 % de ladite uricase sont sous forme octamère.

25

30

35

40


45

50

55

Figure 1

UV Assay of Uricolytic Activity in Fractions from Mono Q Chromatography of PKS Uricase
 (Protein Based on Size-Exclusion HPLC)



**Size-Exclusion HPLC on Superdex 200 of Unfractionated PKS Uricase
(Load) and Mono Q Column Fractions in the Low-Salt Pool**




FIG. 2

Figure 3

Size-Exclusion HPLC on Superdex 200 of Mono Q
Column Fractions of PKS Uricase in the High-Salt Pool




Figure 4



Octamer Content of Mono Q Column Fractions of PKS Uricase

Figure 5


UV Uricase Assays of Sera from Mice Injected with $6 \times 10\text{-kDa PEG}$ Conjugates of PKS Uricase or of Pools from Mono Q Column Fractions
Mice Were Bled 24 Hours after Each Weekly Injection.



Data for the Low-Salt and High-Salt Pools were shifted on the x-axis by 0.1 and 0.2 units, respectively.

Figure 6

**ELISA Analyses with Pig Uricase on the Plate and 1:30 Sera from Mice Injected with
 $6 \times 10\text{-kDa}$ PEG Conjugates of Unfractionated PKS Uricase or Low-Salt or High-Salt Pools**



REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 370084 A [0005] [0018]
- US 9917514 W [0005]
- US 5643575 A [0020]
- WO 0008196 A [0022]
- US 5612460 A [0025]
- US 5286637 A [0025]
- DD 279486 A1 [0025]
- US 5458135 A [0033]
- US 5653974 A [0035]

Non-patent literature cited in the description

- Wu, X et al. *J Mol Evol*, 1992, vol. 34, 78-84 [0004]
- Hande, KR et al. *Am J Med*, 1984, vol. 76, 47-56 [0004]
- Fam, AG. *Baillière's Clin Rheumatol*, 1990, vol. 4, 177-192 [0004]
- Pui, C-H. *Leukemia*, 1997, vol. 11, 1813-1816 [0004]
- Donadio, D et al. *Nouv Presse Méd*, 1981, vol. 10, 711-712 [0004]
- Leaustic, M et al. *Rev Rhum Mal Osteoartic*, 1983, vol. 50, 553-554 [0004]
- Henney et al. *N. Engl. J. Med*, 1968, vol. 278, 2244-2246 [0007]
- Moore et al. *J. Clin Endocrinol. Metab.*, 1980, vol. 51, 691-697 [0007]
- Braun et al. *Pharm. Res.*, 1997, vol. 14, 1472-1478 [0007]
- Braun et al. *Pharm. Res.*, 1997, vol. 14, 1394-1400 [0007]
- PCR Technology. Principles and Applications for DNA Amplification. Stockton Press, 1989 [0023]
- Sambrook, J et al. Molecular Cloning. A laboratory Manual. Cold Spring Harbor Laboratory Press, 1989 [0023]
- Fridovich, I. *J Biol Chem*, 1965, vol. 240, 2491-2494 [0023]
- FM et al. *Appl Biochem Biotechnol*, 1985, vol. 11, 141-152 [0025]
- Moore, WV et al. *J Clin Endocrinol Metab*, 1980, vol. 51, 691-697 [0030]
- Patton, JS. *Adv Drug Delivery Rev*, 1996, vol. 19, 3-36 [0033]
- Remington's Pharmaceutical Sciences. Mack Publishing Co, 1990 [0034]
- Stealth Liposomes. CRC Press, 1995 [0035]
- Venkataseshan, VS et al. *Nephron*, 1990, vol. 56, 317-321 [0036]
- Lee et al. *Science*, 1988, vol. 239, 1288-1291 [0038]
- Wu et al. *Proc. Natl. Acad. Sci. U. S. A.*, 1989, vol. 86, 9412-9416 [0038]
- Veronese, FM et al. *Appl Biochem Biotechnol*, 1985, vol. 11, 141-152 [0041]
- Kito, M et al. *J Clin Biochem Nutr*, 1996, vol. 21, 101-111 [0041]
- Sherman, MR et al. Poly(ethylene glycol) Chemistry and Biological Applications. American Chemical Society, vol. 680, 155-176 [0041]
- Kunitani, M et al. *J Chromatogr*, 1991, vol. 588, 125-137 [0041]
- I. Fridovich. *J Biol Chem.*, 1965, vol. 240, 2491-2494 [0044]
- P. Fossati et al. *J. Clin Chem*, 1980, vol. 26, 227-231 [0046]
- B. Porstmann et al. *J Clin. Chem. Clin. Biochem.*, 1981, vol. 19, 435-440 [0047]